

NEW CIRCUMSCRIPTION AND COMBINATIONS IN THE TRIBE *LEPTOCEREAE* BUXTAUM SENSU GUIGGI (CACTACEAE-CACTOIDEAE) FROM THE CARIBBEAN REGION

Leptocereae Buxbaum *sensu* Guiggi. **Annotarum:** Buxbaum (1958: 178-179) described a new *Tribus* “*Leptocereae*“ [*genus typicum*: *Leptocereus* (A.Berger) Britton et Rose] for the most primitive genera of the subfamily *Cereoideae* K. Schumann (cfr. *Cactoideae* Eaton, Nyffeler & Eggli, 2010: 120-121) including also *Neoabbottia* Britton et Rose and many others presumed archaic genera from South America (e.g. *Armatocereus* Backeberg, *Neoraimondia* Britton et Rose, *Eulychnia* Philippi, etc). In a recent molecular and morphological analysis (Barrios *et al.*, 2020: 5-7, 12) to avoid the paraphyly in *Leptocereus* *sensu lato*, the accepted genus *Dendrocereus* Britton et Rose (Anderson, 2001: 216-217; Hunt *et al.*, 2006: 74) is submerged in it. However, this procedure is not necessary because in the cladograms included in Barrios *et al.* (2020: 6, 8) is possible to recognize a *monophylum* [i.e. *Leptocereae* Buxbaum *sensu* Guiggi] with three different *cladi* or evolutive lines [i.e. *Leptocereus* (A.Berger) Britton et Rose *sensu stricto*, *Dendrocereus* Britton et Rose, *Neoabbottia* Britton et Rose] with the same bio-geographical history in the Caribbean region, that corrisponding to the follow circumscribed valid genera, easily distinguished by the reproductive characters of their flowers and seeds. Furthermore, pending a new phylogenetic analysis, *Serrulatocereus* Guiggi (2018: 1) could also belong to this tribe.

Dendrocereus Britton et Rose [Arborescent to 10 m high; ribs 3-5, to 7 cm high; flower funneliform, solitary, nocturnal, to 20 cm long, tube long, slender, with small, reflexed scales, and few, short spines, perianth segments long-oblong, whitish; fruit hard, naked, to 12 cm long; seeds with testa cells strongly convex at the periphery, ca. 3.6 x 2.8 mm; distributed in Cuba, S Hispaniola; incl. *D. undulosus* (De Candolle) Britton et Rose and subsp. *nudiflorus* (Engelmann ex C.Wright) Guiggi; cfr. Britton & Rose, 1920: 113; Backeberg, 1955: 50-53, 1960: 1948-1950, 1977: 127; Barthlott & Hunt, 2000: 36, 63; Anderson, 2001: 216; Hunt *et al.*, 2006: 74; Barrios *et al.*, 2020: 11].

Leptocereus (A.Berger) Britton et Rose [Normally shrubby or creeping, < 1-3(8) m high; ribs 3-8(-9), to 2 cm high; flower tubular- or obconical-campanulate, solitary or in cluster, diurnal or nocturnal, to 5.5(7) cm long, tube short, stout, spiny, perianth segments short-ovate, reflexed, normally colored; fruit fleshy, normally spiny and ribbed, 1.5-8(-10) cm long; seeds with testa cells flats or verrucous, pitted, ca. 2.5 x 1.6-2 mm; distributed in Cuba, Dominican Republic; incl. *L. arboreus* Britton et Rose, *L. assurgens* (C.Wright ex Grisebach) Britton et Rose (= *Leptocereus ekmanii* (Werdermann) F.M.Knuth, = *L. prostratus* Britton et Rose), *L. assurgens* subsp. *albellus* (Areces) Guiggi, *L. assurgens* subsp. *chrysotyrius* (Areces) Guiggi, *L. carinatus* Areces, *L. demissus* Areces (*holo.* and *iso.* not found, *fide* the JBSD and NY databases; *lectotypus hic designatus: ico. 6*, a branch with an apical cluster of flowers of the type specimen “Areces 6812” in Areces, 2017: 118), *L. leonii* Britton et Rose (= *L. scopulophilus* Areces, = *L. wrightii* León), *L. sylvestris* Britton et Rose (= *L. maxonii* Britton et Rose, = *L. santamarinae* Areces); cfr. Britton & Rose, 1920: 77; Backeberg, 1960: 1958-1964, 1977: 222-223; Anderson, 2001: 391-395; Barthlott & Hunt, 2000: 36, 63; Hunt *et al.*, 2006: 143-144; Areces-Mallea, 1992: 45-46, 1993a: 227-229, 1993b: 245-248, 2017: 118-121, 2018: 258-266; Barrios *et al.*, 2020: 6, 11].

Neoabbottia Britton et Rose [Arborescent to 10 m high or shrubby; ribs normally 4 or 3-7, to 2.5 cm high; flower tubular, in group or solitary from the apical felted areoles, nocturnal, to 7 cm long, tube slender, with longitudinal scales, sometimes with bristles or cluster of spines, perianth segments short-oblong, normally greenish- or yellowish-white; fruit hard, naked or with spines, to 7 cm long; seeds with testa cells tubercled or crested at the periphery, ca. 2.5 x 1.7 mm; distributed in S Hispaniola, Puerto Rico, British Virgin Islands; incl. *N. grantiana* (Britton) Buxbaum, *N. paniculata* (Lamarck) Britton et Rose (= *N. paniculata* var. *humboldtii* Backeberg), *N. quadricostata* (Bello) Guiggi, *N. weingartiana* (E.Hartmann) Guiggi; cfr. Britton & Rose, 1921: 2-3; Backeberg, 1955: 50-53, 1960: 1950-1958, 1977: 322; Barthlott & Hunt, 2000: 36, 63; Anderson, 2001: 392-395; Hunt et al., 2006: 143-144; Areces-Mallea, 2017: 116-118; Barrios et al., 2020: 11].

Combinationes et stati novi

Dendrocereus Britton et Rose, The Cact. 2: 113 (1920).

Dendrocereus undulosus (De Candolle) Britton et Rose **subsp. *nudiflorus*** (Engelmann ex C.Wright) Guiggi **comb. et stat. nov.** *Basionymus*: *Cereus nudiflorus* Engelmann ex C.Wright in Sauvalle, in Anales Acad. Ci. Méd. Fís. Nat. Habana Revista Ci. 6: 98. 1869. **Typus**: Cuba, on the shores, near Habana and Guantánamo, 1866-1867, C.Wright 3570 [MO, holo.?; GH, iso.; NY iso. spec. vis., corp, cos, ar, sp, fl, cfr. Barrios et al., 2020: 12]. *Synonymi*: *Dendrocereus nudiflorus* (Engelmann ex C.Wright) Britton et Rose, The Cact. 2: 113 (1920); *Leptocereus nudiflorus* (Engelmann ex C.Wright) D.Barrios et S.Arias, in Pl. Syst. Evol. 306: 12 (2020). **Distributio**: Cuba. **Annotarum**: a geographical subspecies with a disjunct distribution in Cuba (vs. Hispaniola) characterized by the shorter flowers (to 12 cm long vs. to 20 cm), while the vegetative characters are identicals (cfr. Britton & Rose, 1920: 113, 1925: 221; Hunt et al., 2006: 74).

Leptocereus (A. Berger) Britton et Rose, in Contr. U.S. Natl. Herb. 12(10): 433 (1909).

Leptocereus assurgens (C.Wright ex Grisebach) Britton et Rose **subsp. *albellus*** (Areces) Guiggi **comb. et stat. nov.** *Basionymus*: *Leptocereus assurgens* var. *albellus* Areces, in Cact. Succ. J. (U.S.) 90(4): 260 (2018). **Typus**: Cuba, Prov. Pinar del Río, approximately 20 km NNE from Guane, in Sierra de San Carlos mountain range, on cliffs surrounding the “Hoyo de los Helechos,” a deep solution hole within the Majagua-Cantera cavernous system, 28 Mar. 1992, A.E.Areces 6343 [HAJB holo., NY, iso.; not found cfr. Barrios et al. (2020: 12), Guiggi (2020: 5)]. **Lectotypus (designatus** by Barrios et al. 2020: 12): ico. 5, flower of the type specimen “Areces 6343” in Areces (2018: 260). **Epitypus** (designatus by Barrios et al. 2020: 12): Cuba, Pinar del Río, Minas de Matahambre, mogote de la cueva de Cando, en ladera sur, Sierra de San Carlos, 15 May 2018, D.Barrios et J.L.Guerra HFC90015 [HAJB, epi.; HAJB isoepi.]. *Synonymus*: *Leptocereus albellus* (Areces) Barrios et S. Arias, in Pl. Syst. Evol. 306: 12 (2020). **Distributio**: W Cuba. **Annotarum**: the DNA analysis (Barrios et al. 2020: 6) confirms this taxon as nested in the *Leptocereus assurgens* complex cfr. Hunt et al. (2006: 143-144), with distinguishable vegetative and reproductive characters and a peculiar habitat (cfr. Areces, 2018: 262; Guiggi, 2020: 5), as consequence the recognition as an infra-specific taxon is more appropriate as suggested by Areces (2018: 260).

Leptocereus assurgens (C.Wright ex Griseb.) Britton et Rose **subsp. *chrysotyrius*** (Areces) Guiggi **comb. et stat. nov.** *Basionymus*: *Leptocereus chrysotyrius* Areces, in Cact. Succ. J. (U.S.) 90(4): 263 (2018). **Typus**: Cuba, Prov. Pinar del Río, near the Río San Diego, 1 km N of San Diego de los Baños, 3 Nov. 1976, A.E.Areces 4547 [HAJB holo., NY, iso.; not found fide the HAJB and NY databases]. **Lectotypus hic designatus**: ico. 8, flower of the type specimen “Areces 4547” in Areces (2018: 263). **Distributio**: W Cuba. **Annotarum**: a recognized subspecies distinguished

by the spines reddish-brown, longer, to 53 mm (vs. yellowish, to 2 cm long) and a flower tube purplish (vs. yellowish-green), *cfr.* Areces (2018: 263-265), Hunt. *et al.* (2006: 143-144).

Neoabbotia Britton *et* Rose, in Smithsonian Misc. Collect. 72, no. 9: 2 (1921).

Neoabbotia quadricostata (Bello) Guiggi **comb. nov.** *Basionymus:* *Cereus quadricostatus* Bello, in Anal. Soc. Esp. Hist. Nat. 10: 276 (1881). *Typus:* Puerto Rico, on the shores of Guánica, *D. Bello, non servatus* (*cfr.* Hunt *et al.* 2006: 144). *Neotypus (designatus* by Santiago-Valentín *et al.* 2015: 336): Puerto Rico, *prope* Guayanilla *in montibus calcareis ad los Indios*, 29 Aug. 1886, *P.Sintenis* 4919 *sub Cereus quadricostatus* [B, *neo.*; US, *isoneo. spec. vis.*, corp, cos, ar, sp; K *isoneo. spec. vis.*, corp, cos, ar, sp, fl, fr; BM, G, GH, W]. *Synonymus:* *Leptocereus quadricostatus* (Bello) Britton *et* Rose, in Contr. U.S. Natl. Herb. 16: 242 (1913). **Distributio:** Puerto Rico, British Virgin Islands.

Neoabbotia weingartiana (E.Hartmann) Guiggi **comb. nov.** *Basionymus:* *Cereus weingartianus* E.Hartmann in Dams, in Monatsschr. Kakteenk. 14: 155 (1904). *Typus:* Haiti, *ex cult., non servatus* (*cfr.* Hunt *et al.* 2006: 144). *Lectotypus hic designatus:* ico. in Hartmann, *loc. cit.* 14: 155 (1904). *Epitypus hic designatus:* Haiti, Petite Gonave Island, 9-10 Jul. 1920, *E.C.Leonard* 5256 *sub Leptocereus weingartianus* [NY, *epi. spec. vis.*, corp, cos, ar, sp, fig. 1] *Synonymus:* *Leptocereus weingartianus* (E.Hartmann) Britton *et* Rose, The Cact. 2: 77 (1920). **Distributio:** Hispaniola. **Annotarum:** the neo- and isotypes designated by Areces (2017: 121) have not been found *fide* the JBSD and NY database.

Bibliographia

- Anderson, E. F. 2001. *The Cactus Family*. Timber Press: Portland Oregon.
- Areces-Mallea, A.E. 1992. *Leptocereus santamarinae* (Cactaceae), a new species from Cuba. *Brittonia*. 44(1): 45-49.
- _____. 1993a. *Leptocereus scopulophilus* (Cactaceae), a new species from western Cuba. *Brittonia*. 45(3): 226-230.
- _____. 1993b. *Leptocereus carinatus* (Cactaceae): una nueva especie de la isla de Cuba. *Moscosoa*. 7: 243-248
- _____. 2017. *Leptocereus demissus* a new species from southwestern Hispaniola. *Cact Succ J. (U.S.)*. 89(3): 115-121.
- _____. 2018. New taxa of *Leptocereus* (Cactaceae: Cactoideae) from western Cuba. *Cact. Succ. J. (U.S.)*. 90(4): 258-266.
- Backeberg, C. 1955. *Dendrocereus* and *Neoabbotia*. *Cact. Succ. J. (U.S.)*. 27: 48-53.
- _____. 1960. *Die Cactaceae*. Bd. 4. Gustav Fischer Verlag: Jena (Germany).
- _____. 1977. *Cactus Lexicon*. Blandford Press: Poole, Dorset (England).
- Barrios, D., L.R. González-Torres, S. Arias, and L.C. Majure. 2020. Phylogeny and taxonomy of the Antillean endemic genus *Leptocereus* (Cactaceae) inferred from chloroplast markers and morphological evidence. *Pl. Syst. Evol. (on-line version)*. 306(63): 1-15
- Barthlott, W., and D. Hunt. 2000. Seed-diversity in the Cactaceae subfam. Cactoideae. *Succulent Plant Research*. Vol. 5. DH Books: Milborne Port (England).
- Bello, D. 1881. Apuntes para la flora de Puerto-Rico. Pt. 1. *Anales Soc. Esp. Hist. Nat.* 10: 276-277.
- Britton, N. L., and J. N. Rose. 1920. *The Cactaceae*. Vol. 2. Carnegie Institute: Washington.
- _____. 1921. *Neoabbotia*, a new cactus genus from Hispaniola. *Smithsonian Misc. Collect.* 72(9): 1-6.
- _____. 1925. The tree cactuses of the West Indies. *J. New York Bot. Gard.* 26(310): 217-221.

- Buxbaum, F. 1958. The phylogenetic division of the subfamily *Ceroideae*, *Cactaceae*. *Madroño* 16: 177-216.
- Hartmann, E. 1904. *Cereus weingartianus*. In. Von der Kakteen-Ausstellung Düsseldorf. Dams E. (ed.). *Monatsschr. Kakteenk.* 14: 155-157.
- Hunt, D., N. Taylor, and G. Charles (eds.). 2006. *New Cactus Lexicon*. DH Books: Milborne Port (England).
- Guiggi, A. (2018) *Serrulatocereus* Guiggi. A new proposed genus for the recent re-evaluated *Cereus serruliflorus* Haworth from Haiti (*Cactaceae*). *Cactology V (Suppl. VI)*: 1–3.
- _____. 2020. Caribbean region: Nomenclatural revision in *Cactaceae*. *Cactology V (Suppl. VIII)*: 5–8.
- Nyffeler, R., and U. Eggli. 2010. A farewell to dated ideas and concepts: molecular phylogenetics and a revised suprageneric classification of the family *Cactaceae*. *Schumannia*. 6: 109-149.
- Santiago-Valentín, E., L. Sánchez-Pinto and J. Francisco-Ortega. 2015. Domingo Bello y Espinosa (1817–1884) and the new taxa published in his *Apuntes para la flora de Puerto-Rico*. *Taxon*. 64(2): 323-349.



THE NEW YORK
BOTANICAL GARDEN copyright reserved

0cm 1 2 3 4 5 6 7 8 9 10



SMITHSONIAN INSTITUTION
FROM THE UNITED STATES NATIONAL HERBARIUM

PLANTS OF HAITI
BIOLOGICAL EXPLORATION CONDUCTED BY DR. W. L. ABBOTT

LeptoCereus weingartianus (Hartm.) B.P.
Coral rock, common
Petite Gonave Island

No. 5256 E. C. LEONARD, Collector

July 9 and 10, 1920



NY IMAGED

NEW YORK BOTANICAL GARDEN
00948133

Figure. 1: epitype here designated of *Cereus weingartianus* E.Hartmann [\equiv *Neoabbottia weingartiana* (E.Hartmann) Guiggi], collected in Haiti by E.C. Leonard 5256, slender, multi-stemmed sections, deposited at NY. Photo: © New York Botanical Garden.

Edited & published by Alessandro Guiggi
DISTAV, Polo Botanico, Università degli Studi di Genova
International Cactaceae Research Center (ICRC)
alex.guiggi@libero.it

The texts have been written by Alessandro Guiggi
Illustrations by individual contributors

All right reserved

No parts of this issue may be reproduced in any form, without permission from the Publisher
© Copyright ICRC

ISSN 1971-3010 (paper edition)
ISSN 2421-5457 (online edition)

Printed or digital copies have been deposited at the libraries of Hanbury Gardens, Huntington
Botanical Gardens, Kew Gardens, New York Botanical Garden, Jardin Exotique de Monaco and
Sukkulanten Sammlung Zürich

Nomenclatural novelties proposed in this *supplementum*

Dendrocereus undulosus subsp. *nudiflorus* (Engelmann ex C.Wright) Guiggi comb. et stat.
nov.

Leptocereus assurgens subsp. *albellus* (Areces) Guiggi comb. et stat. nov.

Leptocereus assurgens subsp. *chrysotyrius* (Areces) Guiggi comb. et stat. nov.

Neoabbotia quadricostata (Bello) Guiggi comb. nov.

Neoabbotia weingartiana (E.Hartmann) Guiggi comb. nov.

Acknowledgements

I wish to thank A. Tavera (Jardín Botánico Nacional "Dr. Rafael M. Moscoso") and B. Falcón-Hidalgo (Jardín Botánico Nacional, Universidad de La Habana) for the informations about the dry specimens of A.E. Areces-Mallea deposited respectively at JBSD and HAJB, and E. Santiago-Valentín (University of Puerto Rico) for his article on Domingo Bello y Espinosa.